Exploring the three flavor effects with future superbeams using liquid argon detectors

Sanjib Kumar Agarwalla*
Institute of Physics, Sachivalaya Marg, Sainik School Post, Bhubaneswar 751005, India

Suprabh Prakash† and S. Uma Sankar‡
Department of Physics, Indian Institute of Technology Bombay, Powai, Mumbai 400 076, India

arXiv:1304.3251v1 [hep-ph] 11 Apr 2013

Recent measurement of a moderately large value of θ_{13} signifies an important breakthrough in establishing the standard three flavor oscillation picture of neutrinos. It has opened up exciting possibilities for current and future neutrino oscillation experiments. Current experiments are now poised to probe the impact of full three flavor effects to discover neutrino mass hierarchy, CP violation and octant of θ_{23}. However, they can do so only for favorable ranges of parameters at limited C.L. Hence, future facilities are mandatory to cover the entire parameter space at unprecedented C.L. In their first phase, these experiments will have a very limited reach for CP violation discovery. Therefore, for them, the first step would be to focus on hierarchy and octant measurements. In this letter, we explore the capabilities of LBNE and LBNO, whose large matter effects will enable them to settle these two issues. LBNO has a $\gtrsim 10\sigma$ hierarchy discovery even for the most unfavorable combination of neutrino parameters. The hierarchy reach of LBNE is more limited. In particular, if θ_{23} happens to be in the lower octant, then even a 3σ discrimination would not be possible for LBNE in its first stage, for the most unfavorable values of $\delta_{\rm CP}$. If the current experiments find a hint that the parameter combination is unfavorable, then LBNE should increase their exposure. Regarding the octant, the first stage of LBNO has a 4σ discovery potential and that of LBNE has 3σ.

PACS numbers: 14.60.Pq,14.60.Lm,13.15.+g

INTRODUCTION

The discovery of neutrino mixing and oscillations over the past decade provides firm evidence for new physics beyond the Standard Model, needed to explain nonzero neutrino masses and mixing in the leptonic sector. Recently, the unknown 1-3 lepton mixing angle has been measured precisely by the reactor experiments [1–3]. They have found a moderately large value, not too far from its previous upper bound. This represents a significant milestone towards addressing the remaining fundamental questions, in particular determining the neutrino mass hierarchy and searching for CP violation in the neutrino sector. Another recent and crucial development is the indication of non-maximal 2-3 mixing by the MINOS accelerator experiment [4], leading to the problem of determining the correct octant of θ_{23}. The resolution of the above three issues is possible only by observing the impact of three flavor effects in neutrino oscillations. In this letter, we explore the capabilities of future superbeam experiments towards resolving these unknowns, in light of recent measurements.

Oscillation data are insensitive to the lowest neutrino mass. However, it can be measured in tritium beta decay [5], neutrinoless double beta decay [6] and from the contribution of neutrinos to the energy density of the universe [7]. Very recent data from the Planck experiment have set an upper bound over the sum of all the neutrino mass eigenvalues of $\sum m_i \leq 0.23$ eV at 95% C.L. [8]. But, oscillations experiments are capable of measuring the two independent mass-squared differences between the three neutrino masses: $\Delta_{21} = m_2^2 - m_1^2$ and $\Delta_{31} = m_3^2 - m_1^2$. The smallness of the ratio $\alpha = \Delta_{21}/|\Delta_{31}|$ and that of the 1-3 mixing angle θ_{13}, made it possible, so far, to analyze the data from each neutrino oscillation experiment using an appropriate, effective two flavor oscillation approach. This method has been quite successful in measuring the solar and atmospheric neutrino parameters. The next step must involve probing the full three flavor effects, including the sub-leading ones proportional to α. This task will be undertaken, for the first time, by the current oscillation experiments T2K [9] and NOνA [10].

Δ_{21} is required to be positive by the observed energy dependence of the electron neutrino survival probability in solar neutrinos but Δ_{31} is allowed to be either positive or negative by the present data. Hence, two patterns of neutrino masses are possible: $m_3 \gg m_2 > m_1$, called normal hierarchy (NH) where Δ_{31} is positive and $m_2 > m_1 \gg m_3$, called inverted hierarchy (IH) where Δ_{31} is negative. A third possibility, where the three masses are nearly degenerate with very tiny differences between them, also exists with two sub-cases of Δ_{31} being positive or negative. Leptonic CP violation (LCPV) can be established if CP violating phase $\delta_{\rm CP}$ is shown to differ from 0 and 180°. It was not possible to observe a signal for CP violation in the data so far. Thus, $\delta_{\rm CP}$ can have any value in the range $[-180°, 180°]$. Regarding θ_{23}, all the three global fits [11–13] point to a deviation from maximal mixing (MM) *i.e.* $|0.5 - \sin^2\theta_{23}| \neq 0$. This raises an additional question, "Whether θ_{23} lies in the lower octant (LO: $\theta_{23} < 45°$) or higher octant (HO: $\theta_{23} > 45°$)?".

Settling the issue of neutrino mass hierarchy is very important in order to determine the structure of neutrino mass matrix which in turn can give crucial piece of information towards the theory of neutrino masses and mixing [14]. This is also a key ingredient for neutrinoless double beta decay searches probing the Majorana nature of neutrinos. If $\Delta_{31} < 0$, and yet no neutrinoless double beta decay is observed even in the very far future experiments, that would be a strong evidence that neutrinos are not Majorana particles [15]. Another fundamental issue that needs to be addressed in long baseline experiments is to measure $\delta_{\rm CP}$ and to assess LCPV. This new source of low energy CP violation in neutral lepton sector might be helpful to explain the observed matter anti-matter asymmetry in the universe via leptogenesis [16]. A number of innovative ideas, such as $\mu \leftrightarrow \tau$ symmetry [17], A_4 flavor symmetry [18], and quark-lepton complementarity [19] have been invoked to explain the observed pattern of one small and two large mixing angles in the neutrino sector. Measurements of the precise values of θ_{13} and θ_{23} will reveal the pattern of deviations from these symmetries and will lead to a better understanding of neutrino masses and mixing. In particular, the resolution of θ_{23} octant will severely constrain the patterns of symmetry breaking. With the recent discovery of moderately large value of θ_{13}, these three fundamental measurements fall within our reach.

The combined data from the current experiments T2K and NOνA together can provide a hint at 90% confidence level (C.L.) for hierarchy [20] and at 95% C.L. for octant [21]. Only for a very small set of favorable values of $\delta_{\rm CP}$ can they determine these quantities at $> 99\%$ C.L. Discovery of LCPV is possible only for a small range of $\delta_{\rm CP}$ where CP violation is maximum. Moreover, it is not possible to measure $\delta_{\rm CP}$ without first determining the hierarchy [22]. Hence, the determination of mass hierarchy and octant should be considered as the first step towards the discovery of leptonic CP violation. Future facilities must be developed with the requirements that they should have the capability to determine hierarchy and octant at 3σ level or better for any possible value of $\delta_{\rm CP}$ during the first stage and discover CP violation and measure $\delta_{\rm CP}$ during the second stage.

A study of $\nu_\mu \to \nu_e$ and $\bar{\nu}_\mu \to \bar{\nu}_e$ oscillations at long baseline superbeam experiments is the simplest way to probe three flavor effects, including sub-leading ones. Such a study is capable of achieving all the three objectives mentioned above. In this letter, we consider two future long baseline superbeam neutrino experiments, the details of which are given in the next section. In these experiments, the sensitivity to hierarchy arises due to the interference between the standard oscillation amplitudes and the matter effects arising due to long propagation in earth's matter [23]. This interference increases $\nu_\mu \to \nu_e$ oscillation probability ($P(\nu_\mu \to \nu_e)$) for NH and decreases it for IH. For anti-neutrinos the situation is reversed. The dominant term in $P(\nu_\mu \to \nu_e)$ is driven by matter modified Δ_{31} and is proportional to $\sin^2 2\theta_{13} \sin^2 \theta_{23}$ but the sub-dominant term, suppressed by α, depends on $\delta_{\rm CP}$. Since the hierarchy and $\delta_{\rm CP}$ are both unknown, the interplay of these two terms gives rise to hierarchy-$\delta_{\rm CP}$ degeneracy. If the matter effects are large enough, this degeneracy can be broken completely. This is *not* the case for the current experiments NOνA and T2K, because of which their sensitivity to hierarchy is modest for most of the $\delta_{\rm CP}$ range. There is a similar octant-$\delta_{\rm CP}$ degeneracy also, which limits our ability to determine the correct octant of θ_{23}. This problem can be solved by having substantial data in both ν and $\bar{\nu}$ channels [21]. Both the future facilities we consider are designed such that the matter effects are substantial and they break the hierarchy-$\delta_{\rm CP}$ degeneracy completely. And both are scheduled to have equal ν and $\bar{\nu}$ runs.

Matter effects are proportional to the neutrino energy. Hence we need higher energy beams to generate matter effects large enough to break the hierarchy-$\delta_{\rm CP}$ degeneracy. To be close to the oscillation maximum, a larger baseline also has to be chosen. The neutrino interaction cross-sections are higher at higher energies, so the event rates will be larger. But, at higher energies, the neutral current single π^0 background becomes severe. This can be suppressed effectively by doing a full kinematic reconstruction of all the observed particles, which is possible only with exceptionally sensitive detectors with good particle identification and calorimetry. Liquid argon time projection chambers (LArTPCs) have been demonstrated to have these capabilities, which are the choice of detector for both the future facilities we consider.

EXPERIMENTAL SPECIFICATIONS

The long baseline neutrino experiment (LBNE) [24] is one of the major components of Fermilab's intensity frontier program. It will send a new, high intensity, on-axis neutrino beam towards a 10 kt (in its first phase) LArTPC located at Homestake with a baseline of 1300 km. This facility is designed for initial operation at a proton beam power of 708 kW, with proton energy of 120 GeV that will deliver 6×10^{20} protons on target (p.o.t.) in 230 days per calendar year. In our simulation, we have used the latest fluxes being considered by the collaboration, which have been estimated assuming the smaller decay pipe and the lower horn current compared to the previous studies [25]. We have assumed five years of neutrino run and five years of anti-neutrino run. The detector characteristics have been taken from Table 1 of [26]. To have the LArTPC cross-sections, we have scaled the inclusive charged current (CC) cross sections of water by 1.06 (0.94) for the ν ($\bar{\nu}$) case [27, 28].

The long baseline neutrino oscillation experiment

(LBNO) [29] plans to use an experimental setup where neutrinos produced in a conventional wide-band beam facility at CERN would be observed in a proposed 20 kt (in its first phase) LArTPC housed at the Pyhäsalmi mine in Finland, at a distance of 2290 km. The fluxes have been computed [30] assuming an exposure of 1.5×10^{20} p.o.t. in 200 days per calendar year from the SPS accelerator at 400 GeV with a beam power of 750 kW. For LBNO, we consider five years of neutrino run and five years of anti-neutrino run. We assume the same detector properties as that of LBNE.

In the calculations below, we consider three experimental set-ups: LBNE, LBNO and a possible LBNO configuration with a detector half the mass (10 kt), which we denote as 0.5*LBNO. Scaling down the detector size of LBNO by half makes the exposures of these two experiments very similar. Therefore, considering 0.5*LBNO enables us to make a direct comparison between the inherent properties of the two baselines involved. We have used GLoBES [31, 32] for doing simulations.

PHYSICS WITH BI-EVENTS PLOT

In this section, we discuss the main physics content with the help of bi-events plot shown in fig. 1. Here, we have plotted the number of ν_e appearance events vs. the number of $\bar{\nu}_e$ appearance events, for LBNE and 0.5*LBNO for the four possible combinations of hierarchy and octant. Since δ_{CP} is not known, events are generated for the full range $[-180°, 180°]$, which leads to the ellipses.



FIG. 1: Bi-events (ν_e and $\bar{\nu}_e$ appearance) plots for the four octant-hierarchy combinations and all possible δ_{CP} values. The experiments considered are LBNE and 0.5*LBNO. Here $\sin^2 2\theta_{13} = 0.089$. For LO (HO), $\sin^2\theta_{23} = 0.41(0.59)$.

In calculating the event rates, the following input parameters are used: $\Delta_{21} = 7.5 \times 10^{-5}$ eV^2, $\sin^2\theta_{12} = 0.3$ [13], $\Delta m^2_{\mathrm{eff}} = 2.4 \times 10^{-3}$ eV^2 [1][4], and $\sin^2 2\theta_{13} = 0.089$ [1]. For the lower octant (LO) of θ_{23}, the value $\sin^2\theta_{23} = 0.41$ is chosen and for higher octant (HO), it is taken to be 0.59 [13]. Note that we have plotted the total number of events, whereas the actual analysis will be done based on the spectral information. Nevertheless, contours in this figure contain very important information regarding the physics capabilities of the experiments. This can be summarized in the following way: An experiment can determine both the mass hierarchy and the octant of θ_{23}, if every point on a given ellipse is well separated from every point on each of the other three ellipses. The larger the separation, the better is the confidence with which the above parameters can be determined.

For 0.5*LBNO, the two (LO/HO)-IH ellipses are well separated from the two (LO/HO)-NH ellipses, in their number of neutrino events. Hence, 0.5*LBNO has excellent hierarchy determination capability with just neutrino data. However, only neutrino data alone will not be sufficient to determine the octant in case of IH, because various points on (LO/HO)-IH ellipses have the same number of neutrino events. Likewise, only anti-neutrino data cannot determine the octant in case of NH. Therefore, balanced neutrino and anti-neutrino data is mandatory to make an effective distinction between (LO/HO)-IH ellipses and also between (LO/HO)-NH ellipses.

For LBNE, neutrino data alone can not determine hierarchy because various points on LO-NH and HO-IH ellipses have the same number of neutrino events. Thus anti-neutrino data is needed for the determination of hierarchy also. Even with anti-neutrino data, hierarchy determination can be difficult to achieve, if nature chooses LO and one of the two worst case combinations of hierarchy and δ_{CP} which are (NH, 90°) or (IH, −90°). In such a situation, the neutrino and anti-neutrino events are rather close to each other and it will be very difficult for LBNE to reject the wrong combination. Thus, the hierarchy determination at LBNE will suffer for these two *worst case* scenarios. Regarding octant determination, the capability of LBNE is very similar to that of 0.5*LBNO because the separation between the ellipses, belonging to LO and HO are very similar for the two experiments.

RESULTS

Measurement of mass hierarchy and octant should be considered as a prerequisite for the discovery of leptonic CP violation. In their first phases, both LBNE and LBNO will have very minimal CP violation reach: only for 10-20% of the entire range, at 3σ C.L. In this section, we show the discovery reaches that these experiments will have (in their first phases) for hierarchy and octant .

---

[1] The relation between Δ_{31} and $\Delta m^2_{\mathrm{eff}}$ is taken from [33].

Hierarchy discovery

We first focus on the discovery potential of future facilities to exclude the wrong hierarchy. We consider two true values of $\sin^2\theta_{23}$: 0.41 (LO) and 0.5 (MM). The hierarchy reach would suffer the most for LO, hence we show the results for this conservative choice. [2] This gives us four true combinations of θ_{23}-hierarchy: LO-NH, LO-IH, MM-NH and MM-IH. $\Delta\chi^2$ is calculated for each of these four combinations, assuming the opposite hierarchy to be the test hierarchy. In the fit, we marginalize over test $\sin^2\theta_{23}$ in its 3σ range, Δ_{31} and $\sin^2 2\theta_{13}$ in their 2σ ranges. We considered 5% uncertainty in the matter density, ρ. Priors were added for ρ ($\sigma = 5\%$), $\sin^2 2\theta_{13}$ ($\sigma = 5\%$)[3] and Δ_{31} ($\sigma = 4\%$). $\Delta\chi^2$ is also marginalized over the uncorrelated systematic uncertainties (5% on signal and 5% on background) in the set-ups, so as to obtain a $\Delta\chi^2_{\rm min}$ for every $\delta_{\rm CP}$(true).



FIG. 2: Mass hierarchy discovery as a function of true $\delta_{\rm CP}$ for LBNO, 0.5*LBNO and LBNE. Results are shown for the four possible true θ_{23}-hierarchy combinations. For LO (MM), $\sin^2\theta_{23}({\rm true}) = 0.41$ (0.5). Here $\sin^2 2\theta_{13}({\rm true}) = 0.089$.

Fig. 2 shows the discovery reach for hierarchy as a function of $\delta_{\rm CP}$(true). We see that even 0.5*LBNO has $\gtrsim 10\sigma$ hierarchy discovery potential and for all values of $\delta_{\rm CP}$(true) for all four θ_{23}-hierarchy combinations. The potential of LBNO, of course, is even better. The LBNO baseline is close to bimagic which gives it a particular advantage [34, 35]. LBNO has a particular advantage in hierarchy determination For LBNE, a 5σ discovery of hierarchy is possible for only $\sim 50\%$ of the $\delta_{\rm CP}$(true), irrespective of the true θ_{23}-hierarchy combination. For the unfavorable combinations, *i.e.* NH with $\delta_{\rm CP}$ in the upper half plane or IH with $\delta_{\rm CP}$ in the lower half plane, the hierarchy discovery potential of LBNE suffers. In particular, for LO and the worst case combinations, *LBNE will not be able to provide even a 3σ hierarchy discrimination.* Therefore, LBNE must increase their statistics, if NOνA data indicate that the unfavorable combinations are true. The discovery potential for all three set-ups will be better if θ_{23} happens to lie in HO. But, we checked that, even in such a case, a 5σ discovery is not possible with LBNE for $\sim 30\%$ of the upper half plane of $\delta_{\rm CP}$ for HO-NH true and $\sim 70\%$ of the lower half plane of $\delta_{\rm CP}$ for HO-IH true.

Octant discovery

We next consider the discovery reach of the same setups for excluding the wrong octant. We consider the true values of $\sin^2\theta_{23} = 0.41$(LO) and $\sin^2\theta_{23} = 0.59$(HO), so that we have the following four true combinations of octant and hierarchy: LO-NH, LO-IH, HO-NH and HO-IH. $\Delta\chi^2$ is calculated for each of these four combinations, assuming test $\sin^2\theta_{23}$ values from the other octant. For LO (HO) true, we consider the test $\sin^2\theta_{23}$ range from 0.5 to 0.67 (0.34 to 0.5). Rest of the marginalization procedure (over oscillation parameters as well as systematic uncertainties) is the same as that in the case of hierarchy exclusion except with another difference: the final $\Delta\chi^2$ is marginalized over both the hierarchies as the test hierarchy, to obtain $\Delta\chi^2_{\rm min}$.



FIG. 3: Octant resolving capability as a function of true $\delta_{\rm CP}$ for LBNO, 0.5*LBNO and LBNE. Results are shown for the four possible true octant-hierarchy combinations. For LO (HO), $\sin^2\theta_{23}({\rm true}) = 0.41$ (0.59). Here $\sin^2 2\theta_{13}({\rm true}) = 0.089$.

Fig. 2 shows the discovery reach for octant as a function of δ_{CP}(true). It can be seen that for (LO/HO)-IH true, the sensitivities of LBNE and 0.5*LBNO are quite similar whereas they are somewhat better for 0.5*LBNO if (LO/HO)-NH are the true combinations. For LO-(NH/IH), both LBNE and 0.5*LBNO have more than 3σ discovery of octant while for HO-(NH/IH), the $\Delta\chi^2_{\mathrm{min}}$ varies from ~ 6 to ~ 11. However, with full LBNO, we have more than 3.5σ discovery of octant for all octant-hierarchy combinations. A 5σ discovery of octant is possible only for LO-NH true for $\delta_{\mathrm{CP}}(\text{true}) \in (\sim 20^\circ, \sim 150^\circ)$.

---

[2] If HO is true, then the discovery reaches of these experiments will be better than that for the case of MM; discussed later.

[3] A precision of 5% on $\sin^2 2\theta_{13}$ is expected after the full run of Daya Bay.

CONCLUSIONS

With the recent determination of θ_{13}, a clear and comprehensive picture of the three flavor leptonic mixing matrix has been established. This impressive result has crucial consequences for future theoretical and experimental efforts. It has opened up the possibility to probe the sub-dominant three flavor effects in both current and future long baseline precision oscillation facilities. Recent MINOS data hints at a deviation from maximal 2-3 mixing, causing the octant ambiguity of θ_{23}. In this letter, we explore the physics reach of two future superbeam facilities, LBNE and LBNO, to settle the issues of neutrino mass hierarchy, leptonic CP violation and octant of θ_{23}. Discovery of CP violation and measurement of its phase δ_{CP} at a significant C.L. has to be preceded by a clean measurement of hierarchy and octant. In their first phase, these experiments are better suited for measuring hierarchy and octant compared to CP violation. We find that even a 50% scaled down version of LBNO has a $\sim 10\sigma$ hierarchy discovery potential for all octant-hierarchy combinations and for any δ_{CP} value. However, LBNE suffers in this regard and will not provide a 5σ result for about 50% of the δ_{CP} range. Moreover, it fails to provide even a 3σ hierarchy discovery for LO and the worst case combinations. For octant, both the set-ups have reasonable sensitivities; about 4σ for LBNO and 3σ for LBNE. If the data from NOνA indicates an unfavorable hierarchy-δ_{CP} combination, then LBNE must plan to increase their exposure in the first phase itself so as to have a hierarchy measurement at a high C.L.

Acknowledgement SKA would like to thank M. Bishai, G. Zeller and A. Rubbia for helpful communications. SKA was supported by the DST/INSPIRE Research Grant [IFA-PH-12], Department of Science and Technology, India.

---

[*] Email Address: sanjib@iopb.res.in
[†] Email Address: suprabh@phy.iitb.ac.in
[‡] Email Address: uma@phy.iitb.ac.in

[1] F. An et al. (DAYA-BAY Collaboration), Phys.Rev.Lett. **108**, 171803 (2012), 1203.1669.
[2] J. Ahn et al. (RENO collaboration), Phys.Rev.Lett. **108**, 191802 (2012), 1204.0626.
[3] Y. Abe et al. (Double Chooz Collaboration), Phys.Rev. **D86**, 052008 (2012), 1207.6632.
[4] R. Nichol (MINOS) (2012), talk given at the Neutrino 2012 Conference, June 3-9, 2012, Kyoto, Japan, `http://neu2012.kek.jp/`.
[5] A. Osipowicz et al. (KATRIN Collaboration) (2001), hep-ex/0109033.
[6] I. Avignone, Frank T., S. R. Elliott, and J. Engel, Rev.Mod.Phys. **80**, 481 (2008), 0708.1033.
[7] J. Lesgourgues and S. Pastor, Adv.High Energy Phys. **2012**, 608515 (2012), 1212.6154.
[8] P. Ade et al. (Planck Collaboration) (2013), 1303.5076.
[9] Y. Itow et al. (T2K), pp. 239–248 (2001), hep-ex/0106019.
[10] R. Patterson (NOvA Collaboration) (2012), 1209.0716.
[11] D. Forero, M. Tortola, and J. Valle, Phys.Rev. **D86**, 073012 (2012), 1205.4018.
[12] G. Fogli, E. Lisi, A. Marrone, D. Montanino, A. Palazzo, et al., Phys.Rev. **D86**, 013012 (2012), 1205.5254.
[13] M. Gonzalez-Garcia, M. Maltoni, J. Salvado, and T. Schwetz, JHEP **1212**, 123 (2012), 1209.3023.
[14] C. H. Albright and M.-C. Chen, Phys.Rev. **D74**, 113006 (2006), hep-ph/0608137.
[15] S. Pascoli, S. Petcov, and T. Schwetz, Nucl.Phys. **B734**, 24 (2006), hep-ph/0505226.
[16] M. Fukugita and T. Yanagida, Phys.Lett. **B174**, 45 (1986).
[17] R. N. Mohapatra and S. Nussinov, Phys.Rev. **D60**, 013002 (1999), hep-ph/9809415.
[18] K. Babu, E. Ma, and J. Valle, Phys.Lett. **B552**, 207 (2003), hep-ph/0206292.
[19] H. Minakata and A. Y. Smirnov, Phys.Rev. **D70**, 073009 (2004), hep-ph/0405088.
[20] S. K. Agarwalla, S. Prakash, S. K. Raut, and S. U. Sankar, JHEP **1212**, 075 (2012), 1208.3644.
[21] S. K. Agarwalla, S. Prakash, and S. U. Sankar (2013), 1301.2574.
[22] S. Prakash, S. K. Raut, and S. U. Sankar, Phys.Rev. **D86**, 033012 (2012), 1201.6485.
[23] L. Wolfenstein, Phys. Rev. **D17**, 2369 (1978).
[24] T. Akiri et al. (LBNE Collaboration) (2011), 1110.6249.
[25] Mary Bishai, private communication (2012).
[26] S. K. Agarwalla, T. Li, and A. Rubbia, JHEP **1205**, 154 (2012), 1109.6526.
[27] Geralyn Zeller, private communication (2012).
[28] R. Petti and G. Zeller, *Nuclear Effects in Water vs. Argon*, Tech. Rep. LBNE docdb No. 740.
[29] A. Stahl, C. Wiebusch, A. Guler, M. Kamiscioglu, R. Sever, et al., *Expression of Interest for a very long baseline neutrino oscillation experiment (LBNO)* (2012).
[30] Silvestro di Luise (2012), Poster presented at the ICHEP2012 Conference, July 4-11, 2012, Melbourne, Australia, `www.ichep2012.com.au/`.
[31] P. Huber, M. Lindner, and W. Winter, Comput.Phys.Commun. **167**, 195 (2005), hep-ph/0407333.
[32] P. Huber, J. Kopp, M. Lindner, M. Rolinec, and W. Winter, Comput.Phys.Commun. **177**, 432 (2007), hep-ph/0701187.
[33] H. Nunokawa, S. J. Parke, and R. Zukanovich Funchal, Phys.Rev. **D72**, 013009 (2005), hep-ph/0503283.

[34] S. K. Raut, R. S. Singh, and S. U. Sankar, Phys.Lett. **B696**, 227 (2011), 0908.3741.

[35] A. Dighe, S. Goswami, and S. Ray, Phys.Rev.Lett. **105**, 261802 (2010), 1009.1093.